ADA DENTAL SERVICE RESTRICTIONS IN COVID-19



Updated 4th August

The following system is designed based on published triaging systems in Australia for Dentistry taking into consideration the following key objectives:

- 1) Proportionate, pre-planned response to the possible escalation of COVID-19 based on the evolving community context
- 2) Staged restriction of dental services to reduce transmission risks for COVID-19
- 3) Avoidance of likely burden on medical primary care and emergency services should access to urgent dental care cease.

In all restrictions, <u>urgent dental treatment for</u> people who have been confirmed as COVID-19 positive, should NOT be treated in office-based practice.

Dental treatment for patients identified as moderate or high risk of COVID-19 should be provided under transmission based precautions using appropriate PPE as per ADA Transmission Based Precautions document.

	Services that can be performed	Restricted services, defer treatment
No restrictions	All dental services	No restrictions apply
Level 1 Restrictions	All dental treatments using standard precautions for people who do not meet epidemiological or clinical risk factors for COVID-19 infection transmission.	Defer non-urgent treatment for people who are confirmed, probable or suspected cases in accordance with current Department of Health Case Definitions or State./Territory- specific definitions.
Level 2 Restrictions	Provision of dental treatments that are unlikely to generate aerosols or where aerosols generated have the presence of minimal saliva/blood due to the use of rubber dam. This includes: - Examinations - Simple non-invasive fillings without use of high-speed handpieces - Restorative procedures using high speed handpieces only provided with the use of rubber dam - Non-surgical extractions - Hand scaling (no use of ultrasonic scalers) - Medical management of soft tissue presentations (such as ulcers) - Temporomandibular dysfunction management - Denture procedures - Preventative procedures such as the application of topical remineralising agents e.g. fluoride - Orthodontic treatment	Defer all non-urgent treatments that are likely to generate aerosols which may include the use of - high-speed handpieces without the use of rubber dam - ultrasonic scalers - surgical handpieces. Elective implant dental treatment should be delayed. Defer dental treatment for confirmed, probable or suspected cases of COVID-19 where alternative management options (such as antibiotics or pain control) are appropriate until recovered and released from isolation. Urgent treatment for probable or suspected cases of COVID-19 should be provided under transmission- based precautions, including the use of a fit-checked P2 respirator. Please refer to the ADA Level 2 Decision Tree document. Patients confirmed as COVID-19 positive, should NOT be treated in office-based practice as full airborne precautions are required.

Level 3 Restrictions	Provision of dental treatments	Defer all routine dental treatment not meeting the criteria on the left.
	 extractions analgesics or antimicrobials soft tissue pathology e.g. ulcers 2. where generating aerosols is limited to management of:	Defer dental treatment for confirmed, probable or suspected cases of COVID-19 where alternative management options (such as antibiotics or pain control) are appropriate until recovered and released from isolation. Urgent treatment for probable or suspected cases of COVID-19 should be provided under transmission- based precautions, including the use of a fit-checked P2 respirator. Please refer to the ADA Level 3 Decision Tree document. Patients confirmed as COVID-19 positive, should NOT be treated in office-based practice as full airborne precautions are required.
Level 4 Restrictions	 Significant bleeding Difficulty opening the jaw and/or swallowing Referral from a specialist medical practitioner for assessment or management of a patient receiving urgent medical care for 	Defer all dental treatments for patients not fitting the risk categories identified on the left. Defer dental treatment for confirmed, probable or suspected cases of COVID-19 where alternative management options (such as antibiotics or pain control) are appropriate until recovered and released from isolation. Urgent treatment for probable or suspected cases of COVID-19 should be provided under transmission- based precautions, including the use of a fitchecked P2 respirator. Patients confirmed as COVID-19 positive, should NOT be treated in office-based practice as full airborne precautions are required.
Level 5 Restrictions	No routine dental treatment provided. All patients with acute dental concerns to be directed to emergency care centres.	Any dental treatment without expressed permission from the public health authorities.